

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63

JMB

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®



Induction of Inflammatory Cytokines by Double-stranded and Single-stranded siRNAs is Sequence-dependent and Requires Endosomal Localization

Mouldy Sioud*

Department of Immunology
Molecular Medicine Group
The Norwegian Radium
Hospital, Montebello, N-0310
Oslo, Norway

The potential induction of inflammatory cytokines and interferon responses by small-interfering RNAs (siRNAs) represents a major obstacle for their use as inhibitors of gene expression. Therapeutic applications of siRNAs will require a better understanding of the mechanisms that trigger such unwanted effects, especially in freshly isolated human cells. Surprisingly, the induction of tumor necrosis factor (TNF- α) and interleukin-6 (IL-6) in adherent peripheral blood mononuclear cells (PBMC) was not restricted to double-stranded siRNAs, because induction was also obtained with single-stranded siRNAs (sense or antisense strands). The immunostimulatory effects were sequence-dependent, since only certain sequences are prone to induce inflammatory responses while other not. The induction of TNF- α , IL-6 and interferon α (IFN- α) was chloroquine sensitive and dependent more likely on endosomal Toll-like receptor signaling. Indeed, no significant immunostimulatory effects were detected when either double or single-stranded siRNAs were delivered directly to cytoplasm *via* electroporation. Both RNA types activated a NF- κ B promoter-driven luciferase gene in transiently transfected human adherent PBMC. Moreover, culture of immature dendritic cells with either double or single-stranded siRNAs stimulated interleukin-12 production and induced the expression of CD83, an activation marker. Interestingly, several double-stranded siRNAs did not induced TNF- α , IL-6 and IFN- α production, however, their single-stranded sense or antisense did. Taken together, the present data indicate for the first time that the induction inflammatory cytokines and IFN- α responses by either double-stranded or single-stranded siRNAs in adherent PBMC is sequence-dependent and requires endosomal intracellular signaling. The finding that endosomal localization of self-RNAs (sense strands) can trigger Toll-like receptor signaling in adherent human PBMC is intriguing because it indicates that endosomal self-RNAs can display a molecular pattern capable for activating innate immunity.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: RNA interference; siRNAs; innate immunity; Toll-like receptors; inflammatory cytokines

*Corresponding author

Introduction

RNA interference (RNAi) is an evolutionarily

Abbreviations used: DC, dendritic cells; iDC, immature dendritic cells; siRNA, small-interfering RNA; RNAi, RNA interference; PKR, protein kinase R; TLR, Toll-like receptor; IL, interleukin-; TNF- α , tumor necrosis factor α ; NF- κ B, nuclear factor NF-kappa B; INF, interferon; DOTAP, N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-trimethylammoniummethyl-sulfate.

E-mail address of the author mosioud@ulrik.uio.no

conserved post-transcriptional gene-silencing process, resulting in specific mRNA degradation in several organisms.^{1,2} In this process, long double-stranded (ds) RNAs were processed by the RNase III-like enzyme Dicer to 21–25 nucleotides small-interfering RNAs (siRNAs), which are then incorporated into a protein complex, the RNA-induced silencing complex (RISC). Thereafter, the RISC is remodulated into its active form, which contains the proteins necessary for cleaving the target mRNA at the site, where the guide antisense siRNA strand binds.² RNAi and the related phenomenon of

quelling and post-transcriptional gene silencing (PTGS) have been shown to exist in fungus (*Neurospora*), plants (*Arabidopsis*), invertebrates (*Drosophila* and *C. elegans*) and embryonic mouse and human cells.^{2,3} Notably, the cellular role of RNAi is to maintain the integrity of the genome, to defend cells against viral infection, and to regulate the expression of cellular genes.

In general RNAi was difficult to detect in somatic mammalian cells, since dsRNA structures greater than 30 bp stimulate the IFN pathway, which represents a host response to viral infection where several genes are activated.⁴ Much of the interferon responses are caused by the activation of the dsRNA-dependent protein kinase R (PKR). The antiviral effect of PKR is in part mediated through the phosphorylation of the alpha subunit of the eukaryotic translation initiation factor eIF2 α , which results in the sequestration of the recycling factor eIF-2 β in an inactive complex together with eIF2 α -GDP.⁵ The net effect is global inhibition of protein synthesis. However, it has recently been shown that synthetic 21 nt siRNAs, the effector in RNAi, transfected into human somatic cells can effectively bypass activation of the interferon pathway.⁶ Since then siRNAs have become a powerful tool for genetic analysis and might serve as a potent therapeutic tool for gene silencing.^{7,8} In addition to mRNA cleavage, siRNAs can induce chromatin modification in different organisms. Interestingly, a recent report specified a function for siRNAs in

DNA methylation in mammalian cells.⁹ Contrary to what was thought; recent experiments indicate that exogenously delivered siRNAs can activate the interferon pathway.¹⁰⁻¹⁵ However, there are conflicting reports regarding the extent of off-target effects and interferon induction by siRNA in mammalian cells. For example, Rossi and colleagues noted that none of the three tested chemically made siRNAs induced interferon responses in human cells.¹⁶ These differences could be due to several factors, including differences in the cell types, reagent preparations and siRNA sequences. Notably, in all studies only a few siRNAs were tested using the same experimental conditions. To further address the question of what determines siRNA stimulatory function, a comprehensive analysis of 32 siRNAs targeting different genes was performed. The results indicate that the stimulating capacities of either double-stranded, single-stranded sense or antisense siRNAs are sequence dependent and require endosomal compartments for intracellular signaling.

Results and Discussion

The non-specific effects of siRNAs are sequence-dependent

siRNA were initially thought to be small enough to avoid double-stranded RNA responses.⁶

Table 1. Sequences of the siRNAs used in this study

SiRNAs	Sequence (5'-3') (sense strand)	Target gene
1	GGCCUUCUACCUUCAGACTT	Mouse TNF- α
2	GAUCAUCUUCUAAAAUUCTT	Mouse TNF- α
3	GUUCACCUAGAGCCUAAUAGTT	Human HIF-1
4	CUGAUGACCAGCAACUUGATT	Human HIF-1
5	GACAACCAACUAGUGGUGCTT	Mouse TNF- α
9	GAGGCUGAGACAUAGGCACTT	Mouse TNF- α
7	GAAUCUGAUGACAGGGAGGCTT	Scrambled siRNA
8	GAAGAAGUCGUGCUGCCUUTT	Scrambled siRNA
9	GGUGACAAGAACAUCUCUCCATT	Human neuropilin-1
10	GACCUCAUGUACCACAUUCTT	Human PKC- α
11	GCCAUUGCACUGUGAAUACTT	Mouse Csf-1
12	GUGAUCAUUCAGAGCCAGCTT	Mouse Csf-1 receptor
13	GGCAUCUGGCUUAAGGUGATT	Mouse Csf-1 receptor
14	GACCCUCGAGUCAACAGAGTT	Mouse Csf-1
15	GCAUGCCUUGGAAUUCUUTT	Scrambled siRNA
16	GGCCGAUUGAUCUCAGCGCTT	Human TNF- α
17	GGCCAAAUUUACAUUCUCCTT	Rat NG-2
18	CCAACUAUGACCAGGAUAATT	Human MMP-9
19	GUAGAUCAAUACCCUACACTT	Rat NG-2
20	GGAGCGCACCAUCUUCUUCTT	GFP
21	GACUUGAGCGAGCGCUUUUTT	Human C20orf55
22	GAGAUGAUACCACCGAATT	Human CSPP
23	GCAGAUCUGCGUGCGCCAGTT	Human Frizzled-2
24	GGUUCCAUCGAAUCCUGCATT	Human Wnt-1
25	GAGGCAAUCACCAAUAGCATT	Mouse Basigin
26	GAAGAUUUGCGCAGUGGACTT	Human CSPP
27	GUCCGGGCAGGUCUACUUTT	Mouse TNF- α
28	GGCAUGGAUCUCAAGACATT	Mouse TNF- α
29	CCAACGGCAUGGAUCUCAATT	Mouse TNF- α
30	UGCCCUUCUACAACCAGGATT	Human AKT-1
31	GCUGGAGUACAACUACACTT	GFP
32	GCUGGAGAUCUGAAGAACTT	Human cDNA FLJ34902

253 However, recent studies indicated that they could
 254 induce interferons and cytokine responses.¹⁰⁻¹⁵
 255 Although the reported data provide a note of
 256 caution, the activation of the non-specific pathways
 257 by siRNAs is not well understood. We have
 258 previously examined the response of BALB/c
 259 mice to systemic delivery of siRNAs and found
 260 induction of TNF- α and IL-6 by larger RNAs and
 261 LPS, but not with a chemically synthesized
 262 siRNA.¹⁵ whilst the same siRNA preparation
 263 induced inflammatory cytokine responses in
 264
 265
 266

adherent PBMC *in vitro*.¹⁵ To assess whether the
 siRNA non-specific effects are sequence-dependent
 and to uncover the molecular mechanisms by which
 siRNA activate innate immunity genes, the induc-
 tion of TNF- α and IL-6 secretion by 32 different
 siRNAs was examined in adherent PBMC, an
 enriched monocyte population. Freshly isolated
 cells were treated with the various siRNA
 sequences (see Table 1) for 18 hours. Specific
 ELISA on culture supernatants revealed that
 around 50% of the tested siRNAs induced the
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378





Figure 1. Effects of siRNAs upon TNF- α and IL-6 production in adherent PBMC. Cells were transfected with the indicated siRNA molecules (100 nM) for 18 hours and then TNF- α (a) and IL-6 (b) were measured in the supernatants by ELISA. Results are shown as means of five independent experiments \pm SD. The sequences of the used siRNAs are shown in Table 1. (c) Immunostimulatory siRNA induce TNF- α production in a dose dependent manner. Adherent PBMC were transfected with various concentrations (5–200 nM) of either siRNA 27 for 18 hours. Subsequently, TNF- α was measured in the supernatants by ELISA. All results represent the mean of four or more independent experiments. D, cells were incubated with only DOTAP and transfection buffer.



±SD. (c) siRNA induced TNF-α production in blood monocytes. Purified CD14-positive cells were incubated with either siRNA 27 or 32 (100 nM) for 18 hours, and then TNF-α was measured in the supernatants using ELISA. All results represent the mean of four or more independent experiments

production of TNF-α (Figure 1(a)). Adherent PBMC produced also IL-6 upon stimulation with siRNAs complexed to DOTAP (Figure 1(b)). The profile of IL-6 production is comparable to that of TNF-α. Notably, siRNAs varied dramatically in their ability to induce TNF-α and IL-6 production, suggesting that their activities are sequence-dependent. Only six of the siRNAs examined exhibited a strong immunostimulatory effect, with siRNA-27 being the most effective under our experimental conditions.

To determine the immunostimulatory capacity of siRNAs, cells were transfected with siRNA 27 at a range of concentrations (5–200 nM). After 18 hours in culture, secreted TNF-α was assessed by ELISA (Figure 1(c)). As shown, siRNA 27 stimulated TNF-α production even at low concentrations. Taken together these data indicate that siRNAs can activate the expression of inflammatory cytokines, but their affinity and specificity for a potential cellular target are more likely to be influenced by the siRNA base composition. Sequence alignment of the siRNA sequences revealed no significant homology between the immunostimulatory siRNAs, suggesting the involvement of RNA tertiary structures and/or specific dinucleotides.

Intracellular siRNA delivery is required for cytokine induction

Having found that the non-specific effects of siRNAs are sequence-dependent, I have next investigated whether intracellular siRNA delivery was required. Therefore siRNAs were added directly to the culture medium without complexing them to DOTAP. siRNA 27 activated adherent PMBC to secrete TNF-α when complexed to DOTAP. In contrast, free siRNA did not lead to

379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441

442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504

Figure 2. Characterization of siRNA effects. (a) siRNA induced TNF-α production required liposomal delivery of siRNAs. Adherent PBMC were incubated for 18 hours with either siRNA liposome complexes or naked siRNA 27 (100 nM) in X-VIVO 15 medium, a nuclease free medium. Subsequently, TNF-α was measured in the supernatants by ELISA. Results are shown as means of four independent experiments ±SD. (b) Inhibition of PKR. Adherent PBMC were pretreated, or not, with 2-aminopurine (10 mM) for one hour prior incubation with either siRNA 27 or 32 (100 nM) for 18 hours. Subsequently, TNF-α was measured in the supernatants by ELISA. Results are shown as means of four independent experiments ±SD. (c) siRNA induced TNF-α production in blood monocytes. Purified CD14-positive cells were incubated with either siRNA 27 or 32 (100 nM) for 18 hours, and then TNF-α was measured in the supernatants using ELISA. All results represent the mean of four or more independent experiments

significant cytokine production (Figure 2(a)). A nuclease resistant siRNA 27 gave similar results as its unmodified version (data not shown). These results argue that siRNAs target mainly intracellular compounds. In this respect, a role for PKR in the activation of interferons by siRNAs was recently suggested.¹²

To explore the involvement of the dsRNA recognition protein PKR in this process, the specific inhibitor of PKR, 2-aminopurine, was used. Cells were pretreated with 2-aminopurine (10 mM), a normally used concentration, and subsequently treated with siRNA liposome complexes. Pretreatment with 2-aminopurine reduced, but did not abolish cytokine production by siRNAs (Figure 2(b)). These results would indicate the involvement of other cellular targets in siRNA intracellular signaling.

siRNAs activate TNF-α production in purified human monocytes

Monocytes are essential effector cells in chronic inflammatory disorders and infectious diseases. To perform their function, monocytes need to be activated, either *via* inflammatory cytokines produced by the adaptive immune system or *via* direct stimulation with microbial products. Although a high proportion of the adherent PBMC are monocytes, the capacity of siRNAs to activate TNF-α production in purified blood monocytes was investigated. In these experiments, CD14-positive cell populations were prepared from PBMC by positive selection with immunomagnetic beads coated with anti-CD14 monoclonal antibodies (Dyna, Oslo, Norway), and the cells were subsequently incubated with siRNAs (Figure 2(c)). There




Figure 3. Expression of TLR3 by human monocytes and iDC. (a) Flow cytometry analysis. Freshly isolated monocytes and iDC, monocytes cultured for six days in the presence of GM-CSF and IL-4, were stained either with FITC-labeled anti-TLR3 monoclonal antibody (eBioscience) or control mouse IgG1. After washing, cells were analyzed by flow cytometry. The mean fluorescence of stained cells is shown in the upper-right corner of the individual dot plots. (b) RT-PCR for TLR3 and actin mRNA was performed with monocytes and iDC as described in Experimental Procedures. TLR3 was not detected in human monocytes.

was a significant induction of TNF- α production with the immunostimulatory siRNA 27, whereas no significant activity was seen with the control siRNA 32.

TLR3 is not required for responsiveness to either double-stranded or single-stranded siRNAs in adherent PBMC

Notably, the host sensors that initially detect viral and bacterial antigens and trigger cytokine production has been investigated by several groups,

some of which have indicated the involvement of Toll-like receptors (TLRs). These receptors, which are mammalian homologous of the *Drosophila* Toll, recognize specific structural motifs expressed by microbes.¹⁷⁻¹⁹ So far, ten TLRs have been described in humans, and ligands have been defined for nine of them. TLR1, -2 and -6 are triggered by peptidoglycan and other microbial products, TLR3 by dsRNA, TLR4 by LPS, TLR5 by flagellin, TLR7 and -8 by imidazoquinolines, and TLR9 by unmethylated CpG DNA motifs. Although TLR3 is a receptor for dsRNA and cellular mRNA,^{20,21} it

631 does not contain dsRNA binding motifs. While this
 632 work is underway, a recent study has indicated the
 633 involvement of TLR3 in siRNA signaling and
 634 suggested the involvement of dsRNA-binding
 635 protein that has yet to be identified.¹⁴ In contrast
 636 to macrophages and dendritic cells, however,
 637 human monocytes do not express TLR3 as assessed
 638 by flow cytometry and RT-PCR (Figure 3(a) and (b)).
 639 These results agree with those of a recent study
 640 reporting on the expression of TLR3 by human
 641 monocytes.²² In addition, treatment of either adher-
 642 ent PBMC or purified monocytes with anti-TLR3
 643 monoclonal antibody, known to blocks TLR3
 644 signaling,²³ did not inhibit the production of TNF-
 645 α by immunostimulatory siRNAs (data not shown).


647 **Sense and antisense siRNA strands are highly**
 648 **stimulatory when compared to their double-**
 649 **stranded siRNA counterparts**
 650

651 The data described above indicate that a sub-
 652 group of double-stranded siRNAs can induce the
 653 production of inflammatory cytokines. During this
 654 study, however, I have been surprised by the
 655 finding that both single-stranded sense and anti-
 656 sense siRNAs can trigger the production of pro-
 657 inflammatory cytokines (e.g. TNF- α) and
 658 interferons (IFN- α) in adherent PBMC (Figure 4 as
 659 a representative example). This is in contrast to
 660 what have been reported using various cancer cell
 661 lines.^{12,13} As for the double-stranded siRNAs, the
 662 induction of TNF- α by single-stranded siRNAs was
 663 sequence dependent. In the case of siRNA 27, the

694 sense strand exhibited the strongest effects, whereas
 695 in the case of siRNA 32 the antisense strand exerted
 696 the strongest effect on TNF- α production (Figure
 697 4(a)). Notably, the single-stranded siRNAs were
 698 highly immunostimulatory compared to their
 699 double-stranded siRNA counterparts (e.g. siRNA
 700 32). As shown DOTAP alone did not induce either
 701 TNF- α or IFN- α production, indicating that the
 702 effects are related to the RNA sequences. Direct
 703 addition of either naked double-stranded siRNAs
 704 or single-stranded siRNAs did not affect cytokine
 705 production (data not shown). Moreover, several
 706 double-stranded siRNAs did not induce response,
 707 whereas their corresponding single-stranded sense
 708 or antisense strands did (Figure 4(a), as an
 709 illustration). It is worth noting that the same
 710 single-stranded siRNA preparations were used to
 711 prepare the annealed siRNAs. Taken together, these
 712 observations would indicate that the reported
 713 siRNA effects in adherent PBMC are related to the
 714 siRNA sequences and argue against any possible
 715 contamination with endotoxins.

716 **Cytokine induction by double-stranded siRNAs,**
 717 **sense or antisense siRNA strands requires**
 718 **endosomal acidification**
 719

720 Cationic liposome-delivered siRNAs are
 721 expected to enter the cell *via* endocytosis. Because
 722 several Toll-like receptors, in particular TLR7,
 723 TLR8, TLR9, are localized in the endosomes, I
 724 hypothesized that endocytically introduced RNA
 725 could trigger the activation of these receptors,
 726



727 **Figure 4.** Effects of double-
 728 stranded and single-stranded siRNAs upon TNF- α production in
 729 adherent PBMC. (a) Cells were
 730 transfected with either double-
 731 stranded siRNAs (d), sense
 732 siRNA strands (s) or antisense
 733 siRNA strands (a) for 18 hours.
 734 Subsequently, TNF- α was
 735 measured in the supernatants by
 736 ELISA. All test RNA molecules
 737 were used at 100 nM and com-
 738 plexed with DOTAP at 10 μ g/ml.
 739 (b) Inhibition of cytokine pro-
 740 duction by chloroquine in dose-
 741 dependent manner. Cells were
 742 pre-incubated with chloroquine
 743 for two hours prior transfection
 744 with siRNAs (100 nM) for 18
 745 hours. Subsequently, TNF- α was
 746 measured in the supernatants by
 747 ELISA. (c) The test RNA molecu-
 748 les (100 nM) were delivered to the
 749 cells either *via* DOTAP or electro-
 750 poration as indicated on the figure.
 751

687 After 18 hours transfection time, TNF- α was measured in the supernatants by ELISA. (d) Induction of IFN- α by double-stranded and single-strand siRNAs. The test molecules (100 nM) were delivered to adherent PBMC *via* either DOTAP or electroporation. To test the involvement of endosomes in siRNA induction of interferons, cells were also pretreated for two hours with chloroquine prior to transfection with DOTAP. After transfection (20 hours), IFN- α was measured in the supernatants by ELISA. All results represent the mean of three or more independent experiments.

leading MAPK kinases and NF-κB activation. To test the involvement of endosomes in siRNA signaling, cells were treated with an inhibitor of lysosomal acidification, chloroquine for two hours prior to transfection with either double-stranded or single-stranded siRNAs. TNF-α secretion by adherent PBMC were inhibited in a dose-dependent manner by chloroquine (Figure 4(b)), indicating that the acidification of the endosomes is important for cellular response to either double-stranded or single-stranded siRNAs. A similar inhibition was obtained with Bafilomycin, an additional inhibitor of endosomal acidification (data not shown).

Cationic liposomes such as DOTAP both protect the siRNA and mediate internalization *via* endocytosis. After entry, the siRNA must escape the endolysosomes and enter the RNAi pathway. The requirement for acidification of endosomes in adherent PBMC was further examined *via* electroporation of siRNAs into adherent PBMC. In contrast to DOTAP, the electroporation method open up pores in the cell, so siRNA molecules can enter the cell directly into the cytoplasm. As shown in Figure 4(c) and (d), no TNF-α and IFN-α induction was obtained when either double-stranded or single-stranded siRNAs were delivered *via* electroporation. Therefore, endosome compartments are required for both double-stranded and single-stranded siRNA intracellular signaling and subsequent cytokines and interferon production by adherent PBMC. Notably, pretreatment of PBMC with chloroquine prior to transfection with siRNA abrogated IFN-α production (Figure 4(d)), indicating for the first time the involvement of endosomes in siRNA induction of interferons.

Endosomal siRNA signaling supports the involvement of the Toll-like receptor 8

Toll-like receptors (TLRs) represent a class of pattern recognition receptors (PPRs) that detect microbes or their components.¹⁹ TLRs are predominantly expressed on the cell surface; however, a subset (TLR7, TLR8, TLR9, and in some cases, TLR3) is retained in intracellular compartments. Notably, TLR9, TLR7 and TLR8 require endosomal acidification for signaling.¹⁹ Because the induction of inflammatory cytokines by siRNAs is chloroquine-sensitive, the data would support the involvement of these receptors in siRNA intracellular signaling. The recent identification of single-stranded RNAs as a ligand for human TLR8^{24,25} would support our initial observation¹⁵ and the present elaborated results suggesting the involvement of endosomal TLRs in single-stranded siRNA signaling. This novel observation invites the question of what serves as the endosomal receptors for double-stranded siRNAs. In acidic microenvironment such as the endosomes, double-stranded siRNA might dissociate and generate single-stranded siRNAs for intracellular signaling. In support of this idea is the observation that certain double-stranded siRNAs are very less effective in inducing inflammatory cytokine production than their corresponding single-stranded sense or anti-sense strand counterparts (Figure 4). Depending of the relative internal stability, certain siRNA duplexes may exist in equilibrium with their single strand forms, which can activate TLR8 and/or other endosomal TLRs (Figure 5). Alternatively, double-stranded siRNAs may signal through endosomal TLR3, known to bind double-stranded RNAs.²¹





Figure 5. A schematic diagram illustrating the intracellular siRNA signaling. A complex of either double-strand or single-strand siRNAs or cationic lipids bind to the plasma membrane electrostatically and are internalized *via* endocytosis. Within the endosomes RNA molecules can bind to endosomal Toll-like receptors, in particular TLR8 and 7, which can activate the transcription factor NF-κB, leading to inflammatory cytokine production. TLR7, 8 and 9 use MyD88 as primary adaptor to activate interferon production *via* undefined signaling intermediates. TLR3 uses the adaptor protein TRIF to activate interferon regulatory factor-3 (IRF-3). The activation of Toll receptors may activate PKR directly.^{19,11}

883 Given that single-stranded siRNA are highly
 884 immunostimulatory compared to their correspond-
 885 ing double-stranded siRNAs, during siRNA
 886 annealing it is important to make sure that all
 887 siRNA molecules exist as siRNA duplexes.
 888 Although all double-stranded siRNA preparations
 889 used in this study were checked by PAGE for the
 890 presence of unannealed single-stranded siRNAs,
 891 some of the double-stranded siRNA effects could
 892 originate from tiny traces of free single-stranded
 893 siRNAs. Whatever the nature of double-stranded
 894 siRNA receptors; however, the present data indicate
 895 for the first time that induction of inflammatory
 896 cytokines and interferon- α by either double-
 897 stranded or single-stranded siRNAs requires endo-
 898 somal acidification in freshly isolated PBMC. There-
 899 fore, the development of carriers that deliver siRNA
 900 directly into the cytoplasm should overcome the
 901 problem of inflammatory cytokine induction by
 902 chemically synthesized siRNA, a major obstacle for
 903 systemic administration of siRNA in patients.
 904 Similarly to siRNA liposome complexes, naked
 905 siRNAs delivered *via* other techniques *in vitro* and
 906 *in vivo* are more likely to be taken up by cells *via*
 907 endocytosis.

908 The finding that several siRNA sense strands
 909 (mRNA sequences) can be highly immunostimula-
 910 tory indicates that endosomal Toll-like receptors
 911 can recognize self-RNA. Thus, how can the immune
 912 system distinguish between self and non-self RNAs
 913 (e.g. viral RNAs,) and how to mount and amplify an
 914 immune response against invading RNA in a
 915 specific manner? The present data argue that the
 916 immune system uses the endosomes or lysosomal
 917 compartments as molecular recognition signature
 918 for RNA in general (whether self or non-self).
 919 In contrast to viral RNAs, self-RNAs exist in the
 920 cytoplasm and therefore cannot enter the endoso-
 921 mal compartments; hence in normal situation the
 922 immune system cannot be activated by self-RNAs.
 923 Similar to single-stranded siRNAs, liposomal
 924 delivery of short single-stranded RNA (21 nt)
 925 derived from HIV-1 TAT mRNA or human TNF- α
 926 mRNA also induced TNF- α and IFN- α production
 927 in adherent PBMC (data not shown). Thus, endo-
 928 somal location of RNA seems to function as a
 929 "danger signal" for TLR recognition, and subse-
 930 quent activation of innate immunity genes. The
 931 observation that certain sequences are more stimu-
 932 latory than others would indicate that endosomal
 933 TLRs, in particular TLR8, have preferences for a
 934 particular nucleotides and/or structures that need
 935 further investigation.

936
 937 **siRNA treatment can activate the NF- κ B**
 938 **promoter**

940 The core TLR signaling pathway uses myeloid
 941 differentiation factor 88 (MyD88) as the primary
 942 adaptor protein and results in NF- κ B activation,
 943 cytokine production and expression of costimula-
 944 tory molecules, such as the class II major histocom-
 945 patibility complex. To address the involvement of




946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
Figure 6. Immunostimulatory siRNAs activate the
 965 NF- κ B transcription factor. A NF- κ B-dependent luciferase
 966 reporter gene was co-transfected in human adherent
 967 PBMC with a control plasmid containing the *Renilla*
 968 luciferase gene under the control of the CMV promoter
 969 for 24 hours. Subsequently; cells were treated with either
 970 siRNA 27 or 32. The sense (S) and the antisense strand (A)
 971 of siRNA 27 and 32, respectively, were also tested.
 972 Eighteen hours later, luciferase activity was determined
 973 using Dual-Luciferase Reporter Assay System (Promega).
 974 The data show the relative firefly luciferase activity
 975 normalized to that of *Renilla* luciferase. Results are
 976 shown as means of three independent experiments \pm SD.

977
 978
 979
 980 NF- κ B, promoter activation studies using firefly
 981 luciferase as a reporter gene was performed. Adherent
 982 PBMC were transiently transfected with NF- κ B-luciferase
 983 gene along with plasmid DNA with the *Renilla* luciferase
 984 gene for 24 hours. Subsequently, the cells were either
 985 mock transfected or transfected with a stimulatory
 986 siRNA, and 18 hours later, cells were harvested, and
 987 equal amounts of cell extracts were subjected to luciferase
 988 assay (Figure 6). The data showed that double-
 989 stranded siRNA 27 could induce around 8-fold
 990 increase in NF- κ B activity, whereas 22-fold increase
 991 were obtained with the single-stranded siRNA
 992 (S27), again identifying single-stranded siRNAs as
 993 an important inducer of NF- κ B activation.

994
 995
 996
 997
 998
 999
Single-stranded and double-stranded siRNAs
stimulate monocyte-derived immature dendritic
cells to produce cytokines and up-regulate
costimulatory molecules

1000
 1001
 1002
 1003
 1004
 1005
 1006
 1007
 1008
 1009
 1010
 1011
 1012
 1013
 1014
 1015
 1016
 1017
 1018
 1019
 1020
 1021
 1022
 1023
 1024
 1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059
 1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085
 1086
 1087
 1088
 1089
 1090
 1091
 1092
 1093
 1094
 1095
 1096
 1097
 1098
 1099
 1100
 1101
 1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110
 1111
 1112
 1113
 1114
 1115
 1116
 1117
 1118
 1119
 1120
 1121
 1122
 1123
 1124
 1125
 1126
 1127
 1128
 1129
 1130
 1131
 1132
 1133
 1134
 1135
 1136
 1137
 1138
 1139
 1140
 1141
 1142
 1143
 1144
 1145
 1146
 1147
 1148
 1149
 1150
 1151
 1152
 1153
 1154
 1155
 1156
 1157
 1158
 1159
 1160
 1161
 1162
 1163
 1164
 1165
 1166
 1167
 1168
 1169
 1170
 1171
 1172
 1173
 1174
 1175
 1176
 1177
 1178
 1179
 1180
 1181
 1182
 1183
 1184
 1185
 1186
 1187
 1188
 1189
 1190
 1191
 1192
 1193
 1194
 1195
 1196
 1197
 1198
 1199
 1200



(b) After 48 hours stimulation with siRNAs, IL-12 was measured in the supernatants using ELISA. Results are shown as means of three independent experiments \pm SD.

Figure 7. Immature DC activated with siRNA upregulated the expression of CD83 and secreted IL-12. Immature dendritic cells were culture for 48 hours in the presence of either siRNA 27 or 32 (100 nM) and then stained with FITC-labeled monoclonal antibodies specific for CD83 (a). Cells were also stained with the corresponding isotype control. The mean fluorescence of stained cells is shown in the upper-right corner of the individual dot plots. Results are shown as means of three independent experiments \pm SD.

such as TNF- α and IL-12, which can induce differentiation of T cells into a T helper cell type 1.²⁶ In addition, these stimuli are known to up-regulate certain costimulatory molecules such as CD40, CD80, CD83 and CD86. This process underlies DC maturation, and it strongly potentiates the ability of DC to activate naive T cells.^{26,27} Therefore, the effects of siRNAs upon the maturation of DC were investigated. To generate iDCs, freshly isolated monocytes were cultured for six days in the presence of GM-CSF (50 ng/ml) and IL-4 (100 ng/ml). Subsequently, they were both transfected with siRNA 27 or siRNA 32 and then analyzed by flow cytometry (Figure 7(a)). Adding stimulatory siRNA 27 to the culture medium upregulated the expression of CD83, whereas no significant effects were obtained with the control siRNA 32. Treatment with siRNA 27 also upregulated the expression of CD86 (data not shown). Furthermore, siRNA 27-treatment resulted in IL-12 production (Figure 7(b)). Similar results were obtained with single-stranded siRNA 27. These results suggest that stimulatory siRNAs can enhance immunity by inducing cytokine production and DC maturation. Thus, the use of stimulatory double-stranded or single-stranded siRNAs as a new generation of adjuvants may facilitate the design of effective vaccines.

The ability to enhance or augment the innate immune response clearly represents a potential powerful way to prevent or treat infections as well as a way to develop cancer vaccines. Recently, it has been discovered that unmethylated CpG dinucleotides in particular base contexts are recognized by the immune system as danger signals.²⁸ In animal models, deoxyoligonucleotides with stimulatory CpG motifs have been shown to be of therapeutic value as adjuvants for conventional and therapeutic vaccines against infectious diseases and tumors.²⁹ As shown in the present study, the stimulatory RNAs, in particular single-stranded siRNAs, activated human monocytes and dendritic cells to produce TNF- α and IL-12, two key cytokines in immune regulation. In addition, the induction of

other cytokines such as IL-6 and interferons is an essential response for the clearance of viral infections.³⁰ Natural killer cells are activated by cytokines and chemokines, including type I IFN and IL-12, which are secreted by dendritic cells.³¹ Therefore, treatment of cancer and infectious diseases might benefit from the induction of cytokine and dendritic cell activation by immunostimulatory siRNAs. TLR activators are already used as adjuvants to boost immune responses in vaccines. However, the success of a given vaccine may depend on appropriate activation of TLR. Therefore, single-stranded sense RNA-27 should be explored as an adjuvant for vaccination and immunotherapy. On the other hand, the development of TLR antagonist holds promise as a new class of anti-inflammatory agents. Although further study is required, some of the non-stimulatory RNAs may function as antagonists.

Selection of siRNA against human TNF- α

One of the key cytokines crucial to many biological processes is TNF- α . Low levels of TNF- α produced by monocytes and tissue macrophages under physiological conditions are expected to be involved in maintaining cellular and tissue homeostatic. The production of TNF- α increases in response to microbial infection and tissue injury. Because of its important role in the pathogenesis of a variety of inflammatory and immune diseases, TNF- α has been identified as a key target for pharmacological modulation.³² Therefore, the development of agents that specifically inhibit TNF- α may provide clinician with a valuable alternative to traditional disease-modifying anti-inflammatory drugs. In this respect, we have shown that siRNA-targeting mouse TNF- α can delay the onset of LPS-induced sepsis.³³ Out of 12 recently tested siRNA targeting mouse TNF- α , siRNA 29 (see Table 1), exhibited the greatest protective effect.

To select, siRNAs against human TNF- α , several siRNAs were designed (Table 2) and tested for their inhibitory effect on siRNA 27-induced TNF- α

Table 2. Sequences of siRNAs targeting human TNF- α

SiRNAs	Sequence (5'-3') (Sense strand)
H1	GAGUGACAAGCCUGUAGCCTT
H2	CUCAGCGCUGAGAUCAAUCTT
H3	CUAGUGGUGCCAGCCGAUGTT
H4	GCGUGGAGCUGAGAGAUAAAT

production in adherent PBMC (Figure 8(a)). Some siRNAs were effective in inhibiting TNF- α production, whereas others had no effects, thus suggesting the specificity of the inhibitory effects. When tested alone, the siRNA inhibitors significantly reduced LPS-induced TNF- α production by adherent human PBMC (Figure 8(b)). The utility of this selection assay may be extended to other inflammatory cytokines and type I interferon to select effective siRNAs in human blood cells. It is worth noting that the immunostimulatory effects of siRNAs are more pronounced in human PBMC as compared to other human cell lines. This may depend on the expression level of Toll-like receptors, which is under investigation.

Collectively, the present study underscores for the first time the importance of the endosome compartments in siRNA intracellular signaling in adherent PBMC. The results can easily be explained by the interaction of RNA molecules with endosomal Toll-like receptors and subsequent induction of inflammatory and interferon responses (see Figure 5). The finding that endosomal localization of self RNA can trigger Toll-like receptors signaling in adherent PBMC is intriguing because it indicates




Figure 8. Selection of siRNAs against human TNF- α . (a) Inhibitory effects of TNF- α siRNAs on TNF- α production by the immunostimulatory siRNA 27. Adherent PBMC cells were co transfected with siRNA 27 and various siRNA targeting the human TNF- α (H1-5). Eighteen hours later supernatants were analyzed for TNF- α contents by ELISA. Results are shown as means of three independent experiments \pm SD. (b) Blockage of LPS-induced TNF- α expression by the inhibitory siRNAs. The cells were transfected for 18 hours with either siRNA 32, H2, H3 or H4 (50 nM). Subsequently, cells were stimulated with LPS (20 ng/ml) for eight hours and then supernatants were analyzed for TNF- α contents by ELISA. Results are shown as means of three independent experiments \pm SD.

that endosomal self RNA can display a molecular pattern capable for triggering innate immunity activation.

Experimental Procedures

siRNAs

The siRNAs used in this study correspond to various molecules targeting genes related to other ongoing projects. All siRNAs were chemically synthesized by Eurogentec (Seraing, Belgium), dissolved in water and annealed in the transfection buffer (20 mM Hepes, 150 mM NaCl, pH 7.4) at 20 μ M. Analysis of LPS/endotoxin levels in siRNA stocks was found to be less than 0.01 EU/ml (PyrogenR, CAMBREX). The sequences of the sense strands and the target genes are shown in Table 1.

Preparation of human cells

Human mononuclear cells were prepared from buffy coats by density gradient centrifugation (Lymphoprep, Nycomed Pharm, Oslo), washed, and then resuspended in RPMI 1640 containing 10% heated inactivated fetal calf serum (FCS). Enriched monocyte populations were isolated by plastic adherence. After three hours incubation at 37 $^{\circ}$ C, non-adherent cells were removed by repeated gentle washing with warm medium. More than 75% of the obtained cells by this technique are CD14+ cells. In addition, CD14+ cell population was prepared from PBMC by positive selection with immunomagnetic beads coated with anti-CD14 monoclonal antibodies (Dyna, Oslo, Norway).

Culture of dendritic cells

Enriched population of CD14 positive monocytes were isolated from human PBMC as described above and cultured for five days in RPMI 1640 supplemented with 10% FCS and antibiotics, in the presence of 50 ng/ml GM-CSF and 100 ng/ml IL-4 to obtain monocyte-derived immature dendritic cells (iDC). After five days in culture, the cells were either mock transfected or transfected with either siRNA 27 or 32 for 48 hours, and then cells were stained with FITC-conjugated monoclonal antibodies specific for CD83 (Immunotech). Cells were also stained with FITC-conjugated normal mouse IgG1, an isotype control.

FACS staining and analysis

Briefly, 2×10^5 cells were incubated in the recommended FITC-labeled antibody dilutions for 30 minutes at 4 $^{\circ}$ C in staining buffer (PBS containing 0.5% FCS and 0.1% azide). Cells were also stained with the corresponding Ab isotype controls. Subsequently, the cells were washed twice with the staining buffer and then analyzed on a FACScan using Cell-Quest software (BD Biosciences). Cells were gated in forward/side scatter and the quadrants were set on the relative isotype controls.

Transfection and ELISA

Cells were seeded at 5×10^4 /well/200 μ l in 96-well plates. After an overnight incubation with siRNAs/liposome complexes, culture supernatants were collected and

cytokine contents were assessed by ELISA. Cells were transfected in RPMI 1640 medium (PAA Laboratories GmbH) supplemented with 10% heat-inactivated FCS, 2 mM L-glutamine, 100 µg/ml penicillin and 100 µg/ml streptomycin using DOTAP (10 µg/ml) as described.³³ Each assay was carried out in triplicate. When naked siRNAs were used, cells were incubated in X-VIVO 15 medium without adding FCS (CAMBREX). In the case of monocytes, 2×10⁴ cells were used per well. The inactivation of PKR was carried out as follows: Cells were incubated with 2-aminopurine (10 mM), a specific inhibitor of PKR, for one hour prior to addition of the siRNA/liposome complexes. After an overnight transfection with the tested molecules, the levels of TNF-α, IL-6 and IL-12p70 in the culture supernatants were measured by available ELISA kits according to the manufacturer's instructions (R&D Pharmingen). Samples were run in triplicate. Human IFN-α was measured using ELISA kit (PBL Biomedical Laboratories).

Transfection and reporter assays

An enriched monocyte population (2×10⁶ cells) were seeded onto a 24-well plate and transfected the following day with firefly luciferase gene (1 µg) under the control of the NF-κB-promoter and pRL-TK DNA (1 µg, Promega) encoding the *Renilla* luciferase gene. After 24 hours, the cells were transfected with siRNAs and then cells were harvested 18 hours later. Luciferase activities were measured using the Dual-Luciferase Reporter Assay System according to the manufacturer's instructions (Promega).

RT-PCR analysis

Total RNA was extracted from freshly isolated monocytes and monocytes-derived iDC using total RNA isolation kit (Pharmacia Biotech). DNase I-treated RNA (10 µg) was reverse-transcribed with *NotI*-d(T)18 primer using the first-strand cDNA synthesis Kit (Pharmacia Biotech) according to the manufacturer's instructions. Polymerase chain reaction (PCR) was performed on 3 µl of cDNA in 50 µl of final volume. The TLR3 primers are:

- 5'-ATTGGGTCTGGGAACATTTCTCTTC-3' (forward primer)
- 5'-GTGAGATTTAAACATTCCTCTTCGC-3' (reverse primer)

As a control, β actin mRNA was also amplified using the following primers:

- 5'-ATCTGGCACCACACCTTCTACAATGAGCTGC G-3' (forward primer)
- 5'-CGTCATACTCCTGCTTGGTGATCCACATCTGC-3'

After 35 cycles of amplification (one minute at 92 °C; one minute at 56 °C; one minute at 72 °C, samples were analyzed by electrophoresis on 1.5% agarose gel and visualized by ethidium bromide staining.

Uncited Reference

- 34.

Acknowledgements

I thank Drs Per Ole Iversen and Anne Dybwad for critical reading of the manuscript, and Lise Forfang for excellent technical assistance with flow cytometry. This work was supported by the Norwegian Cancer Society.

References

1. Fire, A., Xu, S., Montgomery, M. K., Kostas, S. A., Driver, S. E. & Mello, C. C. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, **391**, 806–811.
2. Hannon, G. J. (2002). RNA interference. *Nature*, **418**, 244–251.
3. Hamilton, A. J. & Baulcombe, D. C. (1999). A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science*, **286**, 950–952.
4. Sen, G. C. (2001). Viruses and interferons. *Annu. Rev. Microbiol.* **55**, 255–281.
5. Saunders, L. R. & Barber, G. N. (2003). The dsRNA binding protein family: critical roles, diverse cellular functions. *FASEB J.* **17**, 961–983.
6. Elbashir, S. M., Harborth, M. J., Ledecknel, W., Yalcin, A., Weber, K. & Tuschl, T. (2001). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature*, **411**, 494–498.
7. Sioud, M. (2004). Therapeutic siRNAs. *Trends Pharmacol. Sci.* **25**, 22–24.
8. Hannon, G. J. & Rossi, J. J. (2004). Unlocking the potential of the human genome with RNA interference. *Nature*, **431**, 371–378.
9. Kawasaki, H. & Taira, K. (2004). Induction of DNA methylation and gene silencing by short interfering RNAs in human cells. *Nature*, **431**, 211–217.
10. Semizarov, D. *et al.* (2003). Specificity of short interfering RNA determined through gene expression signatures. *Proc. Natl Acad. Sci. USA*, **100**, 6347–6352.
11. Persengiev, S. P., Zhu, X. & Green, M. R. (2004). Nonspecific, concentration-dependent stimulation and repression of mammalian gene expression by small interfering RNAs (siRNAs). *RNA*, **10**, 12–18.
12. Sledz, C. A., Holko, M., de Veer, M. J., Silverman, R. H. & Williams, B. R. G. (2003). Activation of the interferon system by short-interfering RNAs. *Nature Cell. Biol.* **5**, 834–8340.
13. Bridge, A. J., Pebernard, S., Ducaux, A., Nicoulaz, A.-L. & Iggo, R. (2003). Induction of an interferon response by RNAi vectors in mammalian cells. *Nature Gen.* **34**, 263–264.
14. Kariko, K., Bhuyan, P., Capodici, J. & Weissman, D. (2004). Small interfering RNAs mediate sequence-independent gene suppression and induce immune activation by signaling through Toll-like receptor 3. *J. Immunol.* **172**, 6545–6549.
15. Sioud, M. & Sørensen, D. R. (2003). Cationic liposome-mediated delivery of siRNAs in adult mice. *Biochem. Biophys. Res. Commun.* **312**, 1220–1225.
16. Kim, D.-H., Longo, M., Han, Y., Lundberg, P., Cantin, E. & Rossi, J. J. (2004). Interferon induction by siRNAs and ssRNAs synthesized by phage polymerase. *Nature Biotechnol.* **22**, 312–325.
17. Medzhitov, R., Preston-Hurlburt, P. & Janeway, C. A., Jr (1997). A human homologue of drosophila Toll protein signals activation of adaptive immunity. *Nature*, **388**, 394–397.

1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386

1387 18. Aderem, A. & Ulevitch, R. J. (2000). Toll-like receptors
 1388 in the induction of the innate immune response.
 1389 *Nature*, **406**, 782–787.
 1390 19. Armant, M. A. & Fenton, M. J. (2002). Toll-like
 1391 receptors: a family of pattern recognition receptors
 1392 in mammals. *Gen. Biol.* **3**, 3011.
 1393 20. Alexopoulou, L., Holt, A. C., Medzhitov, R. & Flavell,
 1394 R. A. (2001). Recognition of double-stranded RNA
 1395 and activation of NF- kappa B by Toll-like receptor 3.
 1396 *Nature*, **413**, 732–738.
 1397 21. Kariko, K., Ni, H., Capodici, J., Lamphier, M. &
 1398 Weissman, D. (2004). mRNA is an endogenous ligand
 1399 for Toll-like receptor 3. *J. Biol. Chem.* **279**, 12542–12550.
 1400 22. Matsumoto, M., Funami, K., Tanabe, M., Oshiumi, H.,
 1401 Shingai, M., Seto, Y. *et al.* (2003). Subcellular localiza-
 1402 tion of Toll-like receptor 3 in human dendritic cells.
 1403 *J. Immunol.* **171**, 3154–3162.
 1404 23. Matsumoto, M., Kikkawa, S., Kohase, M., Miyake, K.
 1405 & Seya, T. (2002). Establishment of a monoclonal
 1406 antibody against human Toll-like receptor 3 blocks
 1407 double stranded RNA-mediated signaling. *Biochem.*
 1408 *Biophys. Res. Commun.* **293**, 1364–1369.
 1409 24. Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F.,
 1410 Kirschning, C. & Akira, S. (2004). Species-specific
 1411 recognition of single-stranded RNA *via* toll-like
 1412 receptor 7 and 8. *Science*, **303**, 1526–1529.
 1413 25. Diebold, S. S., Kaisho, T., Hemmi, H., Akira, S. &
 1414 Sousa, C. R. (2004). Innate antiviral responses by
 1415 means of TLR7-mediated recognition of single-
 1416 stranded RNA. *Science*, **303**, 1529–1531.
 1417 26. Langenkamp, A., Messi, M., Lanzavecchia, A. &
 1418 Sallusto, F. (2000). Kinetics of dendritic cell activation:
 1419 impact on priming of TH1, TH2 and nonpolarized T
 1420 cells. *Nature Immunol.* **1**, 311–316.
 1421 27. Lutz, M. B. & Schuler, G. (2002). Immature, semi-
 1422 mature and fully mature dendritic cells: which signals
 1423 induce tolerance or immunity? *Trends Immunol.* **23**,
 1424 445–449.
 1425 28. Krieg, A. M. (1996). Lymphocyte activation by CpG
 1426 dinucleotide motifs in prokaryotic DNA. *Trends*
 1427 *Microbiol.* **4**, 73–76.
 1428 29. Krieg, A. M., Love-Homan, L., Yi, A. K. & Harty, J. T.
 1429 (1998). CpG DNA induces sustained IL-12 expression
 1430 *in vivo* and resistance to *Listeria monocytogenes*
 1431 challenge. *J. Immunol.* **161**, 2428–2434.
 1432 30. Netea, M. G., van der Graaf, C., Van der Meer, J. W. M.
 1433 & Kullberg, B. J. (2004). Toll-like receptors and the
 1434 host defense against microbial pathogens: bringing
 1435 specificity to the innate-immune system. *J. Leukoc.*
 1436 *Biol.* **75**, 749–755.
 1437 31. Raulet, D. H. (2004). Interplay of natural killer cells
 1438 and their receptors with the adaptive immune
 1439 response. *Nature Immunol.* **5**, 996–1002.
 1440 32. Tracey, K. J. & Cerami, A. (1994). Tumour necrosis
 1441 factor: a pleiotropic cytokine and therapeutic targets.
 1442 *Ann. Rev. Med.* **45**, 491–503.
 1443 33. Sørensen, R. & Leirdal, M. (2003). Sioud M Gene
 1444 silencing by systemic delivery of synthetic siRNAs in
 1445 adult mice. *J. Mol. Biol.* **327**, 761–766.
 1446 34. Sioud, M. (1994). Interaction between tumor necrosis
 1447 factor α ribozyme and cellular proteins. *J. Mol. Biol.*
 1448 **242**, 619–629.

Edited by J. Karn

(Received 20 December 2004; received in revised form 3 March 2005; accepted 4 March 2005)